

Transition to Zero Cosmological Constant and Phantom Dark Energy as Solutions Involving Change of Orientation of Space-Time Manifold

E. I. Guendelman and A. B. Kaganovich

Physics Department, Ben Gurion University of the Negev, Beer Sheva 84105, Israel

(Dated: October 28, 2008)

The main conclusion of long-standing discussions concerning the role of solutions with degenerate metric ($g \equiv \det(g_{\mu\nu}) = 0$ and even with $g_{\mu\nu} = 0$) was that in the first order formalism they are physically acceptable and must be included in the path integral. In particular, they may describe topology changes and reduction of "metrical dimension" of space-time. The latter implies disappearance of the volume element $\sqrt{-g}d^4x$ of a 4-D space-time in a neighborhood of the point with $g = 0$. We pay attention that besides $\sqrt{-g}$, the 4-D space-time differentiable manifold possesses also a "manifold volume measure" (MVM) described by a 4-form which is sign indefinite and generically independent of the metric. The first order formalism proceeds with originally independent connection and metric structures of the space-time manifold. In this paper we bring up the question whether the first order formalism should be supplemented with degrees of freedom of the space-time differentiable manifold itself, e.g. by means of the MVM. It turns out that adding the MVM degrees of freedom to the action principle in the first order formalism one can realize very interesting dynamics. Such Two Measures Field Theory enables radically new approaches to resolution of the cosmological constant problem. We show that fine tuning free solutions describing a transition to $\Lambda = 0$ state involve oscillations of $g_{\mu\nu}$ and MVM around zero. The latter can be treated as a dynamics involving changes of orientation of the space-time manifold. As we have shown earlier, in realistic scale invariant models (SIM), solutions formulated in the Einstein frame satisfy all existing tests of General Relativity (GR). Here we reveal surprisingly that in SIM, all ground state solutions with $\Lambda \neq 0$ appear to be degenerate either in g_{00} or in MVM. Sign indefiniteness of MVM in a natural way yields a dynamical realization of a phantom cosmology ($w < -1$). It is very important that for all solutions, the metric tensor rewritten in the Einstein frame has regularity properties exactly as in GR. We discuss new physical effects which arise from this theory and in particular strong gravity effect in high energy physics experiments.

PACS numbers: 04.50.Kd, 02.40.Sf, 95.36.+x, 98.80.-k, 25.75.Dw

I. INTRODUCTION: DEGENERATE METRIC, MANIFOLD VOLUME MEASURE AND ORIENTATION OF THE SPACE-TIME MANIFOLD

Solutions with degenerate metric were a subject of a long-standing discussions starting probably with the paper by Einstein and Rosen[1]. In spite of some difficulty interpreting solutions with degenerate metric in classical theory of gravitation, the prevailing view was that they have physical meaning and must be included in the path integral[2],[3],[4]. As it was shown in Refs.[2],[5], in the first order formulation of an appropriately extended general relativity, solutions with $g(x) \equiv \det(g_{\mu\nu}) = 0$ can describe changes of the space-time topology. Similar idea is realized also in the Ashtekar's variables[6],[7]. There are known also classical solutions[8]-[14] with change of the signature of the metric tensor.

The space-time regions with $g(x) = 0$ can be treated as having 'metrical dimension' $D < 4$ (using terminology by Tseytlin[4]).

The simplest solution with $g(x) = 0$ is $g_{\mu\nu} = 0$ while the affine connection is arbitrary (or, in the Einstein-Cartan formulation, the vierbein $e_\mu^a = 0$ and ω_μ^{ab} is arbitrary). Such solutions have been studied by D'Auria and Regge[3], Tseytlin[4]), Witten[15], Horowitz[5], Giddings[16], Bañados[17]; it has been suggested that $g_{\mu\nu} = 0$ should be interpreted as essentially non-classical phase in which diffeomorphism invariance is unbroken and it is realized at high temperature and curvature.

Now we would like to bring up a question: whether the equality $g(x) = 0$ really with a necessity means that the dimension of the space-time manifold in a small neighborhood of the point x may become $D < 4$? At first sight it should be so because the volume element is

$$dV_{(metrical)} = \sqrt{-g}d^4x. \quad (1)$$

Note that the latter is the "metrical" volume element, and the possibility to describe the volume of the space-time manifold in this way appears after the 4-dimensional differentiable manifold M_4 is equipped with the metric structure. For a solution with $g_{\mu\nu} = 0$, the situation with description of the space-time becomes even worse. However, in spite of lack of the metric, the manifold M_4 may still possess a nonzero volume element and have the dimension $D = 4$. The

well known way to realize it consists in the construction of a differential 4-form build for example by means of four differential 1-forms $d\varphi_a$, ($a = 1, 2, 3, 4$): $d\varphi_1 \wedge d\varphi_2 \wedge d\varphi_3 \wedge d\varphi_4$. Each of the 1-forms $d\varphi_a$ may be defined by a scalar field $\varphi_a(x)$. The appropriate volume element of the 4-dimensional differentiable manifold M_4 can be represented in the following way

$$dV_{(manifold)} = 4!d\varphi_1 \wedge d\varphi_2 \wedge d\varphi_3 \wedge d\varphi_4 \equiv \Phi d^4x \quad (2)$$

where

$$\Phi \equiv \varepsilon_{abcd}\varepsilon^{\mu\nu\lambda\sigma}(\partial_\mu\varphi_a)(\partial_\nu\varphi_b)(\partial_\lambda\varphi_c)(\partial_\sigma\varphi_d). \quad (3)$$

is the volume measure independent of $g_{\mu\nu}$ as opposed to the case of the metrical volume measure $\sqrt{-g}$. In order to emphasize the fact that the volume element (2) is metric independent we will call it a *manifold volume element* and the measure Φ - a *manifold volume measure*.

If $\Phi(x) \neq 0$ one can think of four scalar fields $\varphi_a(x)$ as describing a homeomorphism of an open neighborhood of the point x on the 4-dimensional Euclidean space R^4 . However if one allows a dynamical mechanism of metrical dimensional reduction of the space-time by means of degeneracy of the metrical volume measure $\sqrt{-g}$, there is no reason to ignore a possibility of a similar effect permitting degenerate manifold volume measure Φ . The possibility of such (or even stronger, with a sign change of Φ) dynamical effect seems to be here more natural since *the manifold volume measure Φ is sign indefinite* (in Measure Theory, sign indefinite measures are known as signed measures[18]). Note that the metrical and manifold volume measures are not obliged generically to be simultaneously nonzero.

The original idea to use differential forms as describing dynamical degrees of freedom of the space-time differentiable manifold has been developed by Taylor in his attempt[19] to quantize the gravity. Taylor argued that quantum mechanics is not compatible with a Riemannian metric space-time; moreover, in the quantum regime space-time is not even an affine manifold. Only in the classical limit the metric and connection emerge, that one allows then to construct a traditional space-time description. Of course, the transition to the classical limit is described in Ref.[19] rather in the form of a general prescription. Thereupon we would like to pay attention to the additional possibility which was ignored in Ref.[19]. Namely, in the classical limit not only the metric and connection emerge but also some of the differential forms could keep (or restore) certain dynamical effect in the classical limit. In such a case, the traditional space-time description may occur to be incomplete. **Our key idea** is that one of such lost differential forms, the 4-form (2) survives in the classical limit as describing dynamical degrees of freedom of the volume measure of the space-time manifold, and hence can affect the gravity theory on the classical level too[57].

If we add four scalar fields $\varphi_a(x)$ as new variables to a set of usual variables (like metric, connection and matter degrees of freedom) which undergo variations in the action principle[58] then one can expect an effect of gravity and matter on the manifold volume measure Φ and vice versa. We will see later in this paper that in fact such effects exist and in particular classical cosmology solutions of a significant interest exist where Φ vanishes and changes sign.

As is well known, the 4-dimensional differentiable manifold is orientable if it possesses a differential form of degree 4 which is nonzero at every point on the manifold. Therefore two possible signs of the manifold volume measure (3) are associated with two possible orientations of the space-time manifold. The latter means that besides a dimensional reduction and topology changes on the level of the differentiable manifold, the incorporation of the manifold volume measure Φ allows to realize solutions describing *dynamical change of the orientation of the space-time manifold*.

In the light of existence of two volume measures, the simplest way to take into account this fact in the action principle consists in the modification of the action which should now consist of two terms, one with the usual measure $\sqrt{-g}$ and another - with the measure Φ ,

$$S_{mod} = \int (\Phi L_1 + \sqrt{-g}L_2) d^4x, \quad (4)$$

where two Lagrangians L_1 and L_2 coupled with manifold and metrical volume measures appear respectively. According to our previous experience[22]-[34] in Two Measures Field Theory (TMT) we will proceed with an additional basic assumption that, at least on the classical level, the Lagrangians L_1 and L_2 are independent of the scalar fields $\varphi_a(x)$, i.e. the manifold volume measure degrees of freedom enter into TMT only through the manifold volume measure Φ . In such a case, the action (4) possesses an infinite dimensional symmetry

$$\varphi_a \rightarrow \varphi_a + f_a(L_1), \quad (5)$$

where $f_a(L_1)$ are arbitrary functions of L_1 (see details in Ref.[24]). One can hope that this symmetry should prevent emergence of the scalar fields $\varphi_a(x)$ dependence in L_1 and L_2 after quantum effects are taken into account.

Note that Eq.(4) is just a convenient way for presentation of the theory in a general form. In concrete models studied in the present paper, we will see that the action (4) can be always rewritten in an equivalent form where *each term in the action has its own total volume measure and the latter is a linear combination of Φ and $\sqrt{-g}$* .

In the next section it will be shown that the space-time geometry described in terms of the original metric and connection of the underlying action (4) is not generically Riemannian. However by making use a change of variables to the Einstein frame one can represent the resulting equations of motion in the Riemannian (or Einstein-Cartan) space-time.

In our previous investigations we have shown that TMT enables radically new approaches to resolution of the cosmological constant (CC) problem[24],[25],[33] (for an alternative approach see Ref.[35]). Intrinsic features of TMT allow to realize a scalar field dark energy model where all dependence of the scalar field appears as a result of spontaneous breakdown of the dilatation symmetry. Solutions of this model formulated in the Einstein frame satisfy all existing tests of General Relativity (GR)[30],[31],[34]. A new sort of dynamical protection from the initial singularity of the curvature becomes possible[33]. It also allows us to realize a phantom dark energy in the late time universe without explicit introducing phantom scalar field[33].

In contrast to all our previous investigations of TMT, *the purpose of the present paper* is to study the dynamics of the metric $g_{\mu\nu}$ and the manifold volume measure Φ (used in the underlying action (4)) in a number of TMT models. The main attention is concentrated on the analysis of the amazing features of "irregularity" of $g_{\mu\nu}$ and Φ (involving change of orientation of the space-time manifold) in the course of transitions to a ground state and in the phantom dark energy. It is very important to note immediately that in the Einstein frame the metric tensor has regularity properties exactly as in GR. The organization of the paper is the following. In Sec. II we discuss general features of classical dynamics in TMT. In Sec. III we consider the pure gravity model. In Secs. IV and V, in the framework of a simple scalar field model I, we analyze in detail the behavior of two volume measures in the course of transition to the ground state with zero CC. In Sec. VI we study a (generically broken) intrinsic TMT symmetry which is restored in the ground states; the relation of this symmetry restoration to the old CC problem[36] is also analyzed; a discussion of this effect is continued in Sec.VIII. In Sec.VII we shortly present the scalar field model II with spontaneously broken global scale invariance[25] studied in detail in Ref.[33]. In the framework of such class of models, an interesting dynamics of the metric and the manifold volume measure in the course of transition to ground states is analyzed in Sec.VIII. In section IX we reveal that a possibility to realize a phantom dark energy without explicit introducing a phantom scalar field (demonstrated in [33]) has the origin in a sign indefiniteness of the manifold volume measure (3). Finally, since one cannot check directly whether a tiny/zero cosmological constant is fine-tuned or not, in Sec.X we discuss the new physical effects which arise from this theory and in particular a strong gravity effect in high energy physics experiments.

II. CLASSICAL EQUATIONS OF MOTION

Varying the measure fields φ_a , we get $B_a^\mu \partial_\mu L_1 = 0$ where $B_a^\mu = \varepsilon^{\mu\nu\alpha\beta} \varepsilon_{abcd} \partial_\nu \varphi_b \partial_\alpha \varphi_c \partial_\beta \varphi_d$. Since $Det(B_a^\mu) = \frac{4^{-4}}{4!} \Phi^3$ it follows that if $\Phi \neq 0$ the constraint

$$L_1 = sM^4 = const. \quad (6)$$

must be satisfied, where $s = \pm 1$ and M is a constant of integration with the dimension of mass. Variation of the metric $g^{\mu\nu}$ gives

$$\zeta \frac{\partial L_1}{\partial g^{\mu\nu}} + \frac{\partial L_2}{\partial g^{\mu\nu}} - \frac{1}{2} g_{\mu\nu} L_2 = 0, \quad (7)$$

where

$$\zeta \equiv \frac{\Phi}{\sqrt{-g}} \quad (8)$$

is the scalar field build of the scalar densities Φ and $\sqrt{-g}$.

We study models with the Lagrangians of the form

$$L_1 = -\frac{1}{b_g \kappa} R(\Gamma, g) + L_1^m, \quad L_2 = -\frac{1}{\kappa} R(\Gamma, g) + L_2^m \quad (9)$$

where Γ stands for affine connection, $R(\Gamma, g) = g^{\mu\nu} R_{\mu\nu}(\Gamma)$, $R_{\mu\nu}(\Gamma) = R_{\mu\nu\lambda}^\lambda(\Gamma)$ and $R_{\mu\nu\sigma}^\lambda(\Gamma) \equiv \Gamma_{\mu\nu,\sigma}^\lambda + \Gamma_{\alpha\sigma}^\lambda \Gamma_{\mu\nu}^\alpha - (\nu \leftrightarrow \sigma)$. Dimensionless factor b_g^{-1} in front of $R(\Gamma, g)$ in L_1 appears because there is no reason for couplings of the scalar curvature to the measures Φ and $\sqrt{-g}$ to be equal. We choose $b_g > 0$ and $\kappa = 16\pi G$, G is the Newton constant. L_1^m and L_2^m are the matter Lagrangians which can include all possible terms used in regular (with only volume measure $\sqrt{-g}$) field theory models.

Since the measure Φ is sign indefinite, the total volume measure $(\Phi/b_g + \sqrt{-g})$ in the gravitational term $-\kappa^{-1} \int R(\Gamma, g)(\Phi/b_g + \sqrt{-g})d^4x$ is generically also sign indefinite.

Variation of the connection yields the equations we have solved earlier[24]. The result is

$$\Gamma_{\mu\nu}^\lambda = \{\lambda_{\mu\nu}\} + \frac{1}{2}(\delta_\mu^\alpha \sigma_{,\nu} + \delta_\nu^\alpha \sigma_{,\mu} - \sigma_{,\beta} g_{\mu\nu} g^{\alpha\beta}) \quad (10)$$

where $\{\lambda_{\mu\nu}\}$ are the Christoffel's connection coefficients of the metric $g_{\mu\nu}$ and

$$\sigma_{,\mu} \equiv \frac{1}{\zeta + b_g} \zeta_{,\mu}, \quad (11)$$

If $\zeta \neq \text{const.}$ the covariant derivative of $g_{\mu\nu}$ with this connection is nonzero (nonmetricity) and consequently geometry of the space-time with the metric $g_{\mu\nu}$ is generically non-Riemannian. The gravity and matter field equations obtained by means of the first order formalism contain both ζ and its gradient as well. It turns out that at least at the classical level, the measure fields φ_a affect the theory only through the scalar field ζ .

For the class of models (9), the consistency of the constraint (6) and the gravitational equations (7) has the form of the following constraint

$$(\zeta - b_g)(sM^4 - L_1^m) + g^{\mu\nu} \left(\zeta \frac{\partial L_{1m}}{\partial g^{\mu\nu}} + \frac{\partial L_2^m}{\partial g^{\mu\nu}} \right) - 2L_2^m = 0, \quad (12)$$

which determines $\zeta(x)$ (up to the chosen value of the integration constant sM^4) as a local function of matter fields and metric. Note that the geometrical object $\zeta(x)$ does not have its own dynamical equation of motion and its space-time behavior is totally determined by the metric and matter fields dynamics via the constraint (12). Together with this, since ζ enters into all equations of motion, it generically has straightforward effects on dynamics of the matter and gravity through the forms of potentials, variable fermion masses and selfinteractions[22]-[34].

For understanding the structure of TMT it is important to note that TMT (where, as we suppose, the scalar fields φ_a enter only via the measure Φ) is a constrained dynamical system. In fact, the volume measure Φ depends only upon the first derivatives of fields φ_a and this dependence is linear. The fields φ_a do not have their own dynamical equations: they are auxiliary fields. All their dynamical effect is displayed only in the following two ways: a) in generating the constraint (6) (or (12)); b) in the appearance of the scalar field ζ and its gradient in all equations of motion.

III. PURE GRAVITY TMT MODEL

Let us start from the simplest TMT model with action (4) where

$$L_1 = -\frac{1}{b_g \kappa} R(\Gamma, g) - \Lambda_1, \quad L_2 = -\frac{1}{\kappa} R(\Gamma, g) - \Lambda_2 \quad (13)$$

and Λ_1, Λ_2 are constants. Note that $\Lambda_1 = \text{const.}$ cannot have a physical contribution to the field equations (in this model - only gravitational) because $\Phi \Lambda_1$ is a total derivative. Nevertheless we keep Λ_1 to see explicitly how Λ_1 appears in the result. $\Lambda_2/2$ would have a sense of the cosmological constant in the regular, non TMT, theory (i.e. with the only measure $\sqrt{-g}$).

Following the procedure described in Sec.II we obtain the gravitational equations (7) and the constraint (12) in the following form:

$$R_{\mu\nu}(\Gamma) = \frac{\kappa}{2} \frac{b_g \Lambda_2}{\zeta - b_g} g_{\mu\nu} \quad (14)$$

$$\zeta = b_g - \frac{2\Lambda_2}{sM^4 + \Lambda_1} = \text{const.}, \quad (15)$$

where sM^4 is the constant of integration that appears in Eq.(6) and we have assumed that the total volume measure in the gravitational term of the action is nonzero, that is $\Phi/b_g + \sqrt{-g} \neq 0$.

Since $\zeta = \text{const.}$ the connection $\Gamma_{\mu\nu}^\lambda$, Eq.(10), coincides with the Christoffel's connection coefficients of the metric $g_{\mu\nu}$. Therefore in the model under consideration, the space-time with the metric $g_{\mu\nu}$ is (pseudo) Riemannian. It follows from Eqs.(14) and (15) the resulting Einstein equations

$$G_{\mu\nu}(g) = \frac{\kappa}{2}\Lambda g_{\mu\nu}; \quad \Lambda = \frac{b_g}{2}(sM^4 + \Lambda_1) \quad (16)$$

Constancy of $\zeta(x)$ on the mass shell, Eq.(15), means that for the described solution the manifold and metrical volume measures coincide up to a normalization factor. However, this is true only on the mass shell; if we were try to start from this assumption in the underlying action the resulting solution would be different completely.

The model possesses a few interesting features in what it concerns the CC:

(1) The effective CC Λ appears as a constant of integration (as we noticed above, the parameter Λ_1 has no a physical meaning and it can be absorbed by the constant of integration). The effective regular, non TMT, gravity theory provides the same equations if the cosmological constant is added explicitly.

(2) The effective cosmological constant Λ does not depend at all on the CC-like parameter Λ_2 (which should describe a total vacuum energy density including vacuum fluctuations of all matter fields). The latter resembles the situation in the unimodular theory[37],[38].

(3) Note that Λ becomes very small if the integration constant is chosen such that $sM^4 + \Lambda_1$ is very small. The latter is equivalent to a solution with $\Phi/b_g \gg \sqrt{-g}$. In the limit where the metrical volume measure $\sqrt{-g} \rightarrow 0$ while the manifold volume measure Φ remains nonzero, we get $\Lambda \rightarrow 0$. Thus a $\Lambda = 0$ state is realized for a solution which involves a reduction of the metrical dimension to $D^{(g)} < 4$ and at the same time the dimension of the space-time as a differentiable manifold remains $D^{(m)} = 4$.

(4) In the limit where the free parameter $b_g \rightarrow \infty$, the gravitational term in the underlying action (Eqs.(4) and (13)) with coupling to the manifold measure Φ approaches zero; then TMT takes the form of a regular (non TMT) field theory, but the effective cosmological constant Λ becomes infinite. If we wish to reach a very small value of Λ keeping the integration constant arbitrary, one should take the opposite limit where $b_g \ll 1$. Then in the underlying action, the weight of the gravitational term with coupling to the manifold volume measure Φ increases with respect to the regular one with coupling to the metrical measure $\sqrt{-g}$.

The above speculations can be regarded as a strong indication that TMT possesses a potential for resolution of the CC problem. In the next sections we will study this issue in more realistic models.

IV. SCALAR FIELD MODEL I

Let us now study a model including gravity as in Eqs.(9) and a scalar field ϕ . The action has the same structure as in Eq.(4) but it is more convenient to write down it in the following form

$$S_{mod}^{(1)} = \frac{1}{b_g} \int d^4x \left[-\frac{1}{\kappa}(\Phi + b_g\sqrt{-g})R(\Gamma, g) + (\Phi + b_\phi\sqrt{-g})\frac{1}{2}g^{\mu\nu}\phi_{,\mu}\phi_{,\nu} - \Phi V_1(\phi) - \sqrt{-g}V_2(\phi) \right] \quad (17)$$

The appearance of the dimensionless factor b_ϕ is explained by the fact that without fine tuning it is impossible in general to provide the same coupling of the ϕ kinetic term to the measures Φ and $\sqrt{-g}$. $V_1(\phi)$ and $V_2(\phi)$ are potential-like functions; we will see below that the physical potential of the scalar ϕ is a complicated function of $V_1(\phi)$ and $V_2(\phi)$.

The constraint (12) reads now

$$(\zeta - b_g)[sM^4 + V_1(\phi)] + 2V_2(\phi) + b_g\frac{\delta}{2}g^{\alpha\beta}\phi_{,\alpha}\phi_{,\beta} = 0, \quad (18)$$

where

$$\delta = \frac{b_g - b_\phi}{b_g} \quad (19)$$

Since $\zeta \neq \text{const.}$ the connection (10) differs from the connection of the metric $g_{\mu\nu}$. Therefore the space-time with the metric $g_{\mu\nu}$ is non-Riemannian. To see the physical meaning of the model we perform a transition to a new metric

$$\tilde{g}_{\mu\nu} = (\zeta + b_g)g_{\mu\nu}, \quad (20)$$

where the connection $\Gamma_{\mu\nu}^\lambda$ becomes equal to the Christoffel connection coefficients of the metric $\tilde{g}_{\mu\nu}$ and the space-time turns into (pseudo) Riemannian. This is why the set of dynamical variables using the metric $\tilde{g}_{\mu\nu}$ we call the Einstein

frame. One should point out that *the transformation (20) is not a conformal one* since $(\zeta + b_g)$ is sign indefinite. But $\tilde{g}_{\mu\nu}$ is a regular pseudo-Riemannian metric. For the action (17), gravitational equations (7) in the Einstein frame take canonical GR form with the same $\kappa = 16\pi G$

$$G_{\mu\nu}(\tilde{g}_{\alpha\beta}) = \frac{\kappa}{2} T_{\mu\nu}^{eff} \quad (21)$$

Here $G_{\mu\nu}(\tilde{g}_{\alpha\beta})$ is the Einstein tensor in the Riemannian space-time with the metric $\tilde{g}_{\mu\nu}$ and the energy-momentum tensor reads

$$T_{\mu\nu}^{eff} = \frac{\zeta + b_\phi}{\zeta + b_g} (\phi_{,\mu} \phi_{,\nu} - \tilde{g}_{\mu\nu} X) - \tilde{g}_{\mu\nu} \frac{b_g - b_\phi}{(\zeta + b_g)} X + \tilde{g}_{\mu\nu} V_{eff}(\phi; \zeta, M) \quad (22)$$

where

$$X \equiv \frac{1}{2} \tilde{g}^{\alpha\beta} \phi_{,\alpha} \phi_{,\beta} \quad (23)$$

and the function $V_{eff}(\phi; \zeta, M)$ is defined as following:

$$V_{eff}(\phi; \zeta, M) = \frac{b_g [sM^4 + V_1(\phi)] - V_2(\phi)}{(\zeta + b_g)^2}. \quad (24)$$

The scalar ϕ field equation following from Eq.(17) and rewritten in the Einstein frame reads

$$\frac{1}{\sqrt{-\tilde{g}}} \partial_\mu \left[\frac{\zeta + b_\phi}{\zeta + b_g} \sqrt{-\tilde{g}} \tilde{g}^{\mu\nu} \partial_\nu \phi \right] + \frac{\zeta V_1' + V_2'}{(\zeta + b_g)^2} = 0 \quad (25)$$

The scalar field ζ in Eqs.(22)-(25) is determined by means of the consistency equation (12) which in the Einstein frame (20) takes the form

$$(\zeta - b_g)[sM^4 + V_1(\phi)] + 2V_2(\phi) + \delta \cdot b_g(\zeta + b_g)X = 0. \quad (26)$$

V. MANIFOLD MEASURE AND OLD COSMOLOGICAL CONSTANT PROBLEM: COSMOLOGICAL DYNAMICS WITH $|\Phi|/\sqrt{-g} \rightarrow \infty$

It is interesting to see the role of the manifold volume measure in the resolution of the CC problem. We accomplish this now in the framework of the scalar field model I of previous section. The ζ -dependence of $V_{eff}(\phi; \zeta, M)$, Eq.(24), in the form of inverse square like $(\zeta + b_g)^{-2}$ has a key role in the resolution of the old CC problem in TMT. One can show that if quantum corrections to the underlying action generate nonminimal coupling like $\propto R(\Gamma, g)\phi^2$ in both L_1 and L_2 , the general form of the ζ -dependence of V_{eff} remains similar: $V_{eff} \propto (\zeta + f(\phi))^{-2}$, where $f(\phi)$ is a function. The fact that only such type of ζ -dependence emerges in $V_{eff}(\phi; \zeta, M)$, and a ζ -dependence is absent for example in the numerator of $V_{eff}(\phi; \zeta, M)$, is a direct result of our basic assumption that L_1 and L_2 in the action (4) are independent of the manifold measure fields φ_a .

Generically, in the action (17), $b_\phi \neq b_g$ that yields a nonlinear kinetic term (i.e. the k -essence type dynamics) in the Einstein frame[59]. But for purposes of this section it is enough to take a simplified model with $b_\phi = b_g$ (which is in fact a fine tuning) since the nonlinear kinetic term has no qualitative effect on the zero CC problem. In such a case solving the constraint (26) for ζ and substituting into Eqs.(22)-(25) we obtain equations for scalar-gravity system which can be described by the regular GR effective action with the scalar field potential

$$V_{eff}(\phi) = \frac{(sM^4 + V_1)^2}{4[b_g(sM^4 + V_1(\phi)) - V_2(\phi)]} \quad (27)$$

For an arbitrary nonconstant function $V_1(\phi)$ there exist infinitely many values of the integration constant sM^4 such that $V_{eff}(\phi)$ has the *absolute minimum* at some $\phi = \phi_0$ with $V_{eff}(\phi_0) = 0$ (provided $b_g[sM^4 + V_1(\phi)] - V_2(\phi) > 0$). This effect takes place as $sM^4 + V_1(\phi_0) = 0$ **without fine tuning of the parameters and initial conditions**. Note that the choice of the scalar field potential in the GR effective action in a form proportional to a perfect square like emerging in Eq.(27) would mean a fine tuning.

For illustrative purpose let us consider the model with

$$V_1(\phi) = \frac{1}{2} \mu_1^2 \phi^2, \quad V_2(\phi) = V_2^{(0)} + \frac{1}{2} \mu_2^2 \phi^2. \quad (28)$$

Recall that adding a constant to V_1 does not effect equations of motion, while $V_2^{(0)}$ absorbs the bare CC and all possible vacuum contributions. We take negative integration constant, i.e. $s = -1$, and the only restriction on the values of the integration constant M and the parameters is that denominator in (27) is positive[60].

Consider spatially flat FRW universe with the metric in the Einstein frame

$$\tilde{g}_{\mu\nu} = \text{diag}(1, -a^2, -a^2, -a^2), \quad (29)$$

where $a = a(t)$ is the scale factor. Each cosmological solution ends with the transition to a $\Lambda = 0$ state via damping oscillations of the scalar field ϕ towards its absolute minimum ϕ_0 . The appropriate oscillatory regime in the phase plane is presented in Fig. 1.




FIG. 1: Typical phase curve (in the phase plane $(\phi, \dot{\phi})$) of the scalar field ϕ during the transition to $\Lambda = 0$ state. For illustrative purposes the parameters are chosen such that $V_{eff} = (M^2/2b_g)(\phi^2 - M^2)^2/(\phi^2 + 4M^2)$ and $\phi_0 = \pm M$ and $\delta = 0$. In the case without fine tuning of the parameters $b_g \neq b_\phi$, i.e. $\delta \neq 0$, the phase portrait is qualitatively the same.

It follows from the constraint (26) (where we took $\delta = 0$) that $|\zeta| \rightarrow \infty$ as $\phi \rightarrow \phi_0$. More exactly, oscillations of $sM^4 + V_1$ around zero are accompanied with a singular behavior of ζ each time when ϕ crosses ϕ_0

$$\frac{1}{\zeta} \sim sM^4 + V_1(\phi) \rightarrow 0 \quad \text{as} \quad \phi \rightarrow \phi_0 \quad (30)$$

and ζ^{-1} oscillates around zero together with $sM^4 + V_1(\phi)$. Taking into account that the metric in the Einstein frame $\tilde{g}_{\mu\nu}$, Eq.(29), is regular we deduce from Eq.(20) that the metric $g_{\mu\nu}$ used in the underlying action (17) becomes degenerate each time when ϕ crosses ϕ_0

$$g_{00} = \frac{\tilde{g}_{00}}{\zeta + b_g} \sim \frac{1}{\zeta} \rightarrow 0; \quad g_{ii} = \frac{\tilde{g}_{ii}}{\zeta + b_g} \sim -\frac{1}{\zeta} \rightarrow 0 \quad \text{as} \quad \phi \rightarrow \phi_0, \quad (31)$$

where we have taken into account that the energy density approaches zero and therefore for this cosmological solution the scale factor $a(t)$ remains finite in all times t . Therefore

$$\sqrt{-g} \sim \frac{1}{\zeta^2} \rightarrow 0 \quad \text{and} \quad \Phi = \zeta \sqrt{-g} \sim \frac{1}{\zeta} \rightarrow 0 \quad \text{as} \quad \phi \rightarrow \phi_0 \quad (32)$$

The detailed behavior of ζ , the manifold measure Φ and $g_{\mu\nu}$ - components[61] are shown in Fig. 2.

Recall that the manifold volume measure Φ is a signed measure[18] and therefore it is not a surprise that it can change sign. But TMT shows that including the manifold degrees of freedom into the dynamics of the scalar-gravity system we discover an interesting dynamical effect: a transition to zero vacuum energy is accompanied by oscillations of Φ around zero. Similar oscillations[62] simultaneously occur with all components of the metric $g_{\mu\nu}$ used in the underlying action (17).

The measure Φ and the metric $g_{\mu\nu}$ pass zero only in a discrete set of moments in the course of transition to the $\Lambda = 0$ state. Therefore there is no problem with the condition $\Phi \neq 0$ used for the solution (6). Also there is no problem with singularity of $g^{\mu\nu}$ in the underlying action since

$$\lim_{\phi \rightarrow \phi_0} \Phi g^{\mu\nu} = \text{finite} \quad \text{and} \quad \sqrt{-g} g^{\mu\nu} \sim \frac{1}{\zeta} \rightarrow 0 \quad \text{as} \quad \phi \rightarrow \phi_0 \quad (33)$$




FIG. 2: Oscillations of the measure Φ , the original metric $g_{\mu\nu}$ and the r.h.s. of Eq.(41) during the transition to $\Lambda = 0$ state.

The metric in the Einstein frame $\tilde{g}_{\mu\nu}$ is always regular because degeneracy of $g_{\mu\nu}$ is compensated in Eq.(20) by singularity of the ratio of two measures $\zeta \equiv \Phi/\sqrt{-g}$.

VI. RESTORATION OF INTRINSIC TMT SYMMETRY IN THE COURSE OF TRANSITION TO ZERO COSMOLOGICAL CONSTANT STATE

Let us now turn to intrinsic symmetry of TMT which can reveal itself in a model with only the manifold volume measure Φ . Indeed, if in Eq.(9) we take the limit[63] $b_g \rightarrow 0$ and $L_2^m \rightarrow 0$ then Eq.(12) reads

$$L_1^m - g^{\mu\nu} \frac{\partial L_1^m}{\partial g^{\mu\nu}} = sM^4, \quad \text{if } \zeta \neq 0. \quad (34)$$

If in addition L_1^m is homogeneous of degree 1 in $g^{\mu\nu}$ then the integration constant M must be zero. The simplest example of a model for L_1^m satisfying this property is the massless scalar field. In such a case the theory is invariant under transformations in the space of the scalar fields φ_a

$$\varphi_a \rightarrow \varphi'_a = \varphi'_a(\varphi_b) \quad (35)$$

resulting in the transformation of the manifold volume measure Φ

$$\Phi(x) \rightarrow \Phi'(x) = J(x)\Phi(x), \quad J(x) = \text{Det}\left(\frac{\partial \varphi'_a}{\partial \varphi_b}\right) \quad (36)$$

simultaneously with the local transformation of the metric

$$g_{\mu\nu}(x) \rightarrow g'_{\mu\nu}(x) = J(x)g_{\mu\nu}(x). \quad (37)$$

This symmetry was studied in earlier publications[22] where we called it the local Einstein symmetry (LES).

Consider now linear transformations in the space of the scalar fields φ_a

$$\varphi_a \rightarrow \varphi'_a = A_a^b \varphi_b + C_b, \quad a, b = 1, 2, 3, 4 \quad (38)$$

where $A_a^b = \text{constants}$, $C_b = \text{constants}$. Then LES (35)-(37) is reduced to transformations of the global Einstein symmetry (GES) with $J = \det(A_a^b) = \text{const}$. Notice that the Einstein symmetry contains a $\mathbb{Z}_2$ subgroup of the sign inversions when $J = -1$:

$$\Phi \rightarrow -\Phi, \quad g_{\mu\nu} \rightarrow -g_{\mu\nu} \quad (39)$$

LES as well as GES appear to be explicitly broken if L_1^m is not a homogeneous function of degree 1 in $g^{\mu\nu}$, for example as in the model where L_1^m describes a scalar field with a nontrivial potential[64]. The Lagrangian L_2^m generically breaks the Einstein symmetry too. The transformation of GES originated by the infinitesimal linear transformations $\varphi_a(x) \rightarrow \varphi'_a(x) = (1 + \epsilon/4)\varphi_a(x)$, $\epsilon = \text{const.}$, yields the following variation of the action (4) written in the form $S = \int \mathcal{L} d^4x$ where $\mathcal{L} = \Phi L_1 + \sqrt{-g} L_2$:

$$\delta S = \int \left[-\frac{\partial \mathcal{L}}{\partial g^{\mu\nu}} g^{\mu\nu} + L_1 \frac{\partial \Phi}{\partial \varphi_{a,\mu}} \varphi_{a,\mu} \right] \epsilon d^4x. \quad (40)$$

The first term in (40) equals zero on the mass shell giving the gravitational equation (7); recall that we proceed in the first order formalism. Integrating the second term by part, using Eq.(6) and the definition (3) of the measure Φ , we reduce the variation (40) to $\delta S = \epsilon \int \partial_\mu j^\mu d^4x$ where $\partial_\mu j^\mu = sM^4 \Phi$ and $j^\mu = sM^4 B_a^\mu \varphi_a$. In the presence of topological defects with $\Phi = 0$, Eq.(6) does not hold anymore all over space-time, and one should keep L_1 in the definition of the current: $j^\mu = L_1 B_a^\mu \varphi_a$. In Subs.VIII.D we will see how such a situation may be realized.

To present the current conservation in the generally coordinate invariant form one has to use the covariant divergence. However when doing this using the original metric $g_{\mu\nu}$ we encounter the non-metricity. It is much more transparent to use the Einstein frame (20) where the space-time becomes pseudo-Riemannian and the covariant derivative of the metric $\tilde{g}_{\mu\nu}$ equals zero identically. Thus with the definition $j^\mu = \sqrt{-\tilde{g}} J^\mu$, using the definition of ζ in Eq.(7) and the transformation to the Einstein frame (20) we obtain

$$\tilde{\nabla}_\mu J^\mu \equiv \frac{1}{\sqrt{-\tilde{g}}} \partial_\mu \left(\sqrt{-\tilde{g}} J^\mu \right) = sM^4 \frac{\zeta}{(\zeta + b_g)^2} \quad (41)$$

As one should expect, when $L_2 \equiv 0$ and L_1^m is homogeneous of degree 1 in $g^{\mu\nu}$, i.e. in the case of unbroken GES, the current is conserved because in this case the integration constant $M = 0$.

As we have seen in the framework of the scalar field model of Sec.V, the dynamical evolution pushes $|\zeta| \equiv |\Phi|/\sqrt{-g} \rightarrow \infty$ as the gravity+scalar field ϕ -system approaches (without fine tuning) the $\Lambda = 0$ ground state $\phi = \phi_0$. Therefore according to Eq.(41),

$$\tilde{\nabla}_\mu J^\mu \rightarrow 0 \quad \text{as} \quad \phi \rightarrow \phi_0. \quad (42)$$

For the model of Sec.V, the damping oscillations of the r.h.s. of Eq.(41) around zero are shown in Fig. 2. Thus, the *GES explicitly broken in the underlying action, emerges in the vacuum which, as it turns out, has zero energy density. And vice versa, emergence of GES due to $|\zeta| \rightarrow \infty$ implies, according to Eq.(24), a transition to a $\Lambda = 0$ ground state.*

Other way to reach the same conclusion is to look at the underlying action (17). In virtue of Eq.(33), it is evident that in the course of transition to the ground state, the terms in (17) coupled to the metric volume measure $\sqrt{-g}$ become negligible in comparison with the corresponding terms coupled to the manifold volume measure Φ ; besides the term $-\int V_1(\phi) \Phi d^4x$ (which also breaks the GES) disappears as $\phi \rightarrow \phi_0$. The only terms surviving in the transition to the $\Lambda = 0$ ground state are the following

$$\frac{1}{b_g} \int \Phi d^4x \left[-\frac{1}{\kappa} R(\Gamma, g) + \frac{1}{2} g^{\mu\nu} \phi_{,\mu} \phi_{,\nu} \right] \quad (43)$$

and they respect the GES.

One should notice however that one can regard the GES as the symmetry responsible for a zero CC only if TMT is taken in the strict framework formulated in Sec.I. In fact, let us consider for example a modified TMT model where the manifold volume measure degrees of freedom enter in the Lagrangian L_1 in contrast to our additional basic assumption made Sec.I (after Eq.(4)). Namely let us assume that the Lagrangian L_1 in Eq.(4) involves a term proportional to $\Phi/\sqrt{-g}$ that explicitly breaks the infinite dimensional symmetry (5). To be more concrete we consider a model with the action

$$S = S_{mod}^{(1)} - \lambda \int \frac{\Phi^2}{\sqrt{-g}} d^4x \quad (44)$$

where $S_{mod}^{(1)}$ is the action defined in Eq.(17). Such an addition to the action (17) respects the GES but it is easy to see that it affects the theory in such a way that without fine tuning it is impossible generically to reach a zero CC (see Appendix B).

VII. SCALAR FIELD MODEL II. GLOBAL SCALE INVARIANCE

Let us now turn to the analyze of the results of the TMT model possessing a global scale invariance studied early in detail[25]-[27],[29]-[33]. The scalar field ϕ playing the role of a model of dark energy appears here as a dilaton, and a spontaneous breakdown of the scale symmetry results directly from the presence of the manifold volume measure Φ . In other words, this SSB is an intrinsic feature of TMT.

In the context of the present paper this model is of significant interest because cosmological solutions of the FRW universe exhibit two unexpected results: (a) the ground state as well as the asymptotic of quintessence-like evolution (in co-moving frame) possess certain degeneracies in Φ or $g_{\mu\nu}$; (b) superaccelerating expansion of the universe (phantom cosmology) appears as the direct dynamical effect when $\Phi < 0$, i.e. as the orientation of the space-time manifold is opposite to the regular one. In this section we present the model and some of its relevant results. Regimes (a) and (b) will be analyzed in the next two sections.

The action of the model reads

$$S = \frac{1}{b_g} \int d^4x e^{\alpha\phi/M_p} \left[-\frac{1}{\kappa} R(\Gamma, g)(\Phi + b_g \sqrt{-g}) + (\Phi + b_\phi \sqrt{-g}) \frac{1}{2} g^{\mu\nu} \phi_{,\mu} \phi_{,\nu} - e^{\alpha\phi/M_p} (\Phi V_1 + \sqrt{-g} V_2) \right] \quad (45)$$

and it is invariant under the global scale transformations ($\theta = \text{const.}$):

$$g_{\mu\nu} \rightarrow e^\theta g_{\mu\nu}, \quad \Gamma_{\alpha\beta}^\mu \rightarrow \Gamma_{\alpha\beta}^\mu, \quad \varphi_a \rightarrow \lambda_{ab} \varphi_b \quad \text{where} \quad \det(\lambda_{ab}) = e^{2\theta}, \quad \phi \rightarrow \phi - \frac{M_p}{\alpha} \theta. \quad (46)$$

The appearance of the dimensionless parameters b_g and b_ϕ is explained by the same reasons we mentioned after Eqs.(9) and (17). In contrast to the model of Sec.IV, now we deal with exponential (pre-) potentials where V_1 and V_2 are constant dimensionfull parameters. The remarkable feature of this TMT model is that Eq.(6), being the solution of the equation of motion resulting from variation of the manifold volume measure degrees of freedom, breaks spontaneously the scale symmetry (46): this happens due to the appearance of a dimensionfull integration constant sM^4 in Eq.(6). One can show[33] that in the case of the negative integration constant ($s = -1$) and $V_1 > 0$, the ground state appears to be again (as it was in the scalar field model I of Sec.IV) a zero CC state without fine tuning of the parameters and initial conditions. The behavior of Φ and $g_{\mu\nu}$ in the course of transition to the $\Lambda = 0$ state is qualitatively the same as we observed in Sec.V for the scalar field model I. Therefore in the present paper studying the model (45) we restrict ourself with the choice $s = +1$ and $V_1 > 0$.

Similar to the model of Sec.IV, equations of motion resulting from the action (45) are noncanonical and the space-time is non Riemannian when using the original set of variables. This is because all the equations of motion and the solution for the connection coefficients include terms proportional to $\partial_\mu \zeta$. However, when working with the new metric (ϕ remains the same)

$$\tilde{g}_{\mu\nu} = e^{\alpha\phi/M_p} (\zeta + b_g) g_{\mu\nu}, \quad (47)$$

which we call the Einstein frame, the connection becomes Riemannian and general form of all equations becomes canonical. Since $\tilde{g}_{\mu\nu}$ is invariant under the scale transformations (46), spontaneous breaking of the scale symmetry is reduced in the Einstein frame to the *spontaneous breakdown of the shift symmetry*

$$\phi \rightarrow \phi + \text{const.} \quad (48)$$

After the change of variables to the Einstein frame (47) the gravitational equation takes the standard GR form with the same Newton constant as in the action (45)

$$G_{\mu\nu}(\tilde{g}_{\alpha\beta}) = \frac{\kappa}{2} T_{\mu\nu}^{eff} \quad (49)$$

where $G_{\mu\nu}(\tilde{g}_{\alpha\beta})$ is the Einstein tensor in the Riemannian space-time with the metric $\tilde{g}_{\mu\nu}$. The energy-momentum tensor $T_{\mu\nu}^{eff}$ reads

$$T_{\mu\nu}^{eff} = \frac{\zeta + b_\phi}{\zeta + b_g} \left(\phi_{,\mu} \phi_{,\nu} - \frac{1}{2} \tilde{g}_{\mu\nu} \tilde{g}^{\alpha\beta} \phi_{,\alpha} \phi_{,\beta} \right) - \tilde{g}_{\mu\nu} \frac{b_g - b_\phi}{2(\zeta + b_g)} \tilde{g}^{\alpha\beta} \phi_{,\alpha} \phi_{,\beta} + \tilde{g}_{\mu\nu} V_{eff}(\phi, \zeta; M) \quad (50)$$

where the function $V_{eff}(\phi, \zeta; M)$ is defined as following:

$$V_{eff}(\phi, \zeta; M) = \frac{b_g [M^4 e^{-2\alpha\phi/M_p} + V_1] - V_2}{(\zeta + b_g)^2}. \quad (51)$$

Note that the ζ -dependence of $V_{eff}(\phi, \zeta; M)$ is the same as in Eq.(24) of the model of Sec.IV.

The scalar field ζ is determined by means of the constraint similar to Eq.(26) of Sec.IV

$$(b_g - \zeta) \left[M^4 e^{-2\alpha\phi/M_p} + V_1 \right] - 2V_2 - \delta \cdot b_g(\zeta + b_g)X = 0 \quad (52)$$

where

$$X \equiv \frac{1}{2} \tilde{g}^{\alpha\beta} \phi_{,\alpha} \phi_{,\beta} \quad \text{and} \quad \delta = \frac{b_g - b_\phi}{b_g} \quad (53)$$

The dilaton ϕ field equation in the Einstein frame is reduced to the following

$$\frac{1}{\sqrt{-\tilde{g}}} \partial_\mu \left[\frac{\zeta + b_\phi}{\zeta + b_g} \sqrt{-\tilde{g}} \tilde{g}^{\mu\nu} \partial_\nu \phi \right] - \frac{2\alpha\zeta}{(\zeta + b_g)^2 M_p} M^4 e^{-2\alpha\phi/M_p} = 0. \quad (54)$$

where again ζ is a solution of the constraint (52). Note that the dilaton ϕ dependence in all equations of motion in the Einstein frame appears *only* in the form $M^4 e^{-2\alpha\phi/M_p}$, i.e. it results only from the spontaneous breakdown of the scale symmetry (46).

The effective energy-momentum tensor (50) can be represented in a form of that of a perfect fluid

$$T_{\mu\nu}^{eff} = (\rho + p)u_\mu u_\nu - p\tilde{g}_{\mu\nu}, \quad \text{where} \quad u_\mu = \frac{\phi_{,\mu}}{(2X)^{1/2}} \quad (55)$$

with the following energy and pressure densities resulting from Eqs.(50) and (51) after inserting the solution $\zeta = \zeta(\phi, X; M)$ of Eq.(52):

$$\rho(\phi, X; M) = X + \frac{(M^4 e^{-2\alpha\phi/M_p} + V_1)^2 - 2\delta b_g(M^4 e^{-2\alpha\phi/M_p} + V_1)X - 3\delta^2 b_g^2 X^2}{4[b_g(M^4 e^{-2\alpha\phi/M_p} + V_1) - V_2]}, \quad (56)$$

$$p(\phi, X; M) = X - \frac{(M^4 e^{-2\alpha\phi/M_p} + V_1 + \delta b_g X)^2}{4[b_g(M^4 e^{-2\alpha\phi/M_p} + V_1) - V_2]}. \quad (57)$$

In a spatially flat FRW universe with the metric $\tilde{g}_{\mu\nu} = \text{diag}(1, -a^2, -a^2, -a^2)$ filled with the homogeneous scalar field $\phi(t)$, the ϕ field equation of motion takes the form

$$Q_1 \ddot{\phi} + 3H Q_2 \dot{\phi} - \frac{\alpha}{M_p} Q_3 M^4 e^{-2\alpha\phi/M_p} = 0 \quad (58)$$

where H is the Hubble parameter and we have used the following notations

$$\dot{\phi} \equiv \frac{d\phi}{dt} \quad (59)$$

$$Q_1 = 2[b_g(M^4 e^{-2\alpha\phi/M_p} + V_1) - V_2]\rho_{,X} = (b_g + b_\phi)(M^4 e^{-2\alpha\phi/M_p} + V_1) - 2V_2 - 3\delta^2 b_g^2 X \quad (60)$$

$$Q_2 = 2[b_g(M^4 e^{-2\alpha\phi/M_p} + V_1) - V_2]p_{,X} = (b_g + b_\phi)(M^4 e^{-2\alpha\phi/M_p} + V_1) - 2V_2 - \delta^2 b_g^2 X \quad (61)$$

$$Q_3 = \frac{1}{[b_g(M^4 e^{-2\alpha\phi/M_p} + V_1) - V_2]} \left[(M^4 e^{-2\alpha\phi/M_p} + V_1)[b_g(M^4 e^{-2\alpha\phi/M_p} + V_1) - 2V_2] + 2\delta b_g V_2 X + 3\delta^2 b_g^3 X^2 \right] \quad (62)$$

The non-linear X -dependence appears here in the framework of the fundamental theory without exotic terms in the Lagrangians L_1 and L_2 . This effect follows just from the fact that there are no reasons to choose the parameters b_g and b_ϕ in the action (45) to be equal in general; on the contrary, the choice $b_g = b_\phi$ would be a fine tuning. Thus the above equations represent an explicit example of k -essence[39] resulting from first principles. The system of equations (21), (56)-(58) accompanied with the functions (60)-(62) and written in the metric $\tilde{g}_{\mu\nu} = \text{diag}(1, -a^2, -a^2, -a^2)$ can be obtained from the k -essence type effective action

$$S_{eff} = \int \sqrt{-\tilde{g}} d^4x \left[-\frac{1}{\kappa} R(\tilde{g}) + p(\phi, X; M) \right], \quad (63)$$

where $p(\phi, X; M)$ is given by Eq.(57). In contrast to the simplified models studied in literature[39], it is impossible here to represent $p(\phi, X; M)$ in a factorizable form like $\tilde{K}(\phi)\tilde{p}(X)$. The scalar field effective Lagrangian, Eq.(57), can be represented in the form

$$p(\phi, X; M) = K(\phi)X + L(\phi)X^2 - U(\phi) \quad (64)$$

where the potential

$$U(\phi) = \frac{[V_1 + M^4 e^{-2\alpha\phi/M_p}]^2}{4[b_g(V_1 + M^4 e^{-2\alpha\phi/M_p}) - V_2]} \quad (65)$$

and $K(\phi)$ and $L(\phi)$ depend on ϕ only via $M^4 e^{-2\alpha\phi/M_p}$. Notice that $U(\phi) > 0$ for any ϕ provided

$$b_g > 0, \quad V_1 > 0 \quad \text{and} \quad b_g V_1 \geq V_2, \quad (66)$$

that we will assume in what follows. Note that besides the presence of the effective potential $U(\phi)$, the Lagrangian $p(\phi, X; M)$ differs from that of Ref.[40] by the sign of $L(\phi)$: in our case $L(\phi) < 0$ provided the conditions (66). This result cannot be removed by a choice of the parameters of the underlying action (45) while in Ref.[40] the positivity of $L(\phi)$ was an essential *assumption*. This difference plays a crucial role for a possibility of a dynamical protection from the initial singularity of the curvature studied in detail in Ref[33].

The model allows a power law inflation (where the dilaton ϕ plays the role of the inflaton) with a graceful exit to a zero or tiny cosmological constant state. In what it concerns to primordial perturbations of ϕ and their evolution, there are no difference with the usual (i.e. one-measure) model with the action (63)-(65).

In the model under consideration, the conservation law corresponding to the GES (38) has the form

$$\tilde{\nabla}_\mu J^\mu \equiv \frac{1}{\sqrt{-\tilde{g}}} \partial_\mu \left(\sqrt{-\tilde{g}} J^\mu \right) = s M^4 \frac{\zeta}{(\zeta + b_g)^2} e^{-2\alpha\phi/M_p} \quad (67)$$

with the same definition of the current J^μ as in Sec.VI.

VIII. DEGENERACIES OF g_{00} AND Φ IN $\Lambda \neq 0$ GROUND STATES

A. Fine-Tuned $\delta = 0$ Models

We are going now to analyze some of the cosmological solutions for the late universe in the framework of the scale invariant model of the previous section. These solutions surprisingly exhibit that asymptotically, as $t \rightarrow \infty$, either $g_{00} \rightarrow 0$ or $\Phi \rightarrow 0$.

In the late universe, the kinetic energy $X \rightarrow 0$. Therefore in many cases the role of the nonlinear X dependence becomes qualitatively unessential. This is why, for simplicity, in this section we can restrict ourself with the fine tuned model with $\delta = 0$. In such a case the constraint (52) yields

$$\zeta = \frac{b_g V_1 - 2V_2 + b_g M^4 e^{-2\alpha\phi/M_p}}{V_1 + M^4 e^{-2\alpha\phi/M_p}}, \quad (68)$$

The energy density and pressure take then the form

$$\rho^{(0)} = \rho|_{\delta=0} = \frac{1}{2}\dot{\phi}^2 + U(\phi); \quad p^{(0)} = p|_{\delta=0} = \frac{1}{2}\dot{\phi}^2 - U(\phi), \quad (69)$$

where $U(\phi)$ is determined by Eq.(65). The ϕ -equation (58) is reduced to

$$\ddot{\phi} + 3H\dot{\phi} + \frac{dU(\phi)}{d\phi} = 0. \quad (70)$$

Applying this model to of the late time cosmology of the spatially flat universe and assuming that the scalar field $\phi \rightarrow \infty$ as $t \rightarrow \infty$, it is convenient to rewrite the potential $U(\phi)$ in the form

$$U(\phi) = \Lambda + V(\phi), \quad (71)$$

where

$$\Lambda = \frac{V_1^2}{4(b_g V_1 - V_2)}. \quad (72)$$

is the positive cosmological constant and

$$V(\phi) = \frac{(b_g V_1 - 2V_2)V_1 M^4 e^{-2\alpha\phi/M_p} + (b_g V_1 - V_2)M^8 e^{-4\alpha\phi/M_p}}{4(b_g V_1 - V_2)[b_g(V_1 + M^4 e^{-2\alpha\phi/M_p}) - V_2]}. \quad (73)$$

It is evident that if $b_g V_1 > 2V_2$ or $b_g V_1 = 2V_2$ then $V(\phi)$ is a sort of a quintessence-like potential and therefore quintessence-like scenarios can be realized. This means that the dynamics of the late time universe is governed by the dark energy which consists of both the cosmological constant and the potential slow decaying to zero as $\phi \rightarrow \infty$. In the opposite case, $b_g V_1 < 2V_2$, the potential $V(\phi)$, and also $U(\phi)$, has an absolute minimum at some finite value of ϕ , and therefore the cosmological scenario is different from the quintessence-like. Details of the cosmological evolution starting from the early inflation and up to the late time universe governed by the potential $U(\phi)$ have been studied in Ref.[33] for each of these three cases. Here we want to analyze what kind of degeneracy appears in ground state depending on the region in the parameter space.

B. The case $b_g V_1 > 2V_2$

Let us consider the case when the relation between the parameters V_1 and V_2 satisfies the condition $b_g V_1 > 2V_2$. It follows from Eq.(68) that

$$\zeta \rightarrow \frac{b_g V_1 - 2V_2}{V_1} = \text{const} > 0 \quad \text{as} \quad \phi \rightarrow \infty \quad (74)$$

By making use the (00) component of Eq.(47), we see that

$$g_{00} = \frac{e^{-\alpha\phi/M_p}}{\zeta + b_g} \rightarrow 0 \quad (75)$$

In order to get the asymptotic time dependence of g_{00} and the spatial components of the metric

$$g_{ii} = -\frac{e^{-\alpha\phi/M_p}}{\zeta + b_g} a(t)^2, \quad i = 1, 2, 3 \quad (76)$$

as $t \rightarrow \infty$, we have to know a solution $a = a(t)$, $\phi = \phi(t)$. We can find analytically the asymptotic (as $\phi \rightarrow \infty$) behavior of a cosmological solution for a particular value of the parameter $\alpha = \sqrt{3/8}$. In such a case, keeping only the leading contribution of the ϕ -exponent in Eq.(73), we deal with the following system of equations

$$\left(\frac{\dot{a}}{a}\right)^2 = \frac{1}{3M_p^2} \rho^{(0)} \quad (77)$$

$$\ddot{\phi} + 3\frac{\dot{a}}{a}\dot{\phi} - \frac{2\alpha}{M_p} \frac{V_1(b_g V_1 - 2V_2)}{4(b_g V_1 - V_2)^2} M^4 e^{-2\alpha\phi/M_p} = 0, \quad (78)$$

where $\rho^{(0)}$ is determined by Eq.(69). The exact analytic solution for these equations is as follows[24]:

$$\phi(t) = \text{const.} + \frac{M_p}{2\alpha} \ln(M_p t), \quad a(t) \propto t^{1/3} e^{\lambda t}, \quad \lambda = \frac{1}{M_p} \sqrt{\frac{\Lambda}{3}} \quad (79)$$

where Λ is determined by Eq.(72). Therefore we obtain for the asymptotic cosmic time behavior of the components of the metric $g_{\mu\nu}$

$$g_{00} \sim \frac{1}{t^{1/2}} \rightarrow 0; \quad g_{ii} \sim -t^{1/6} e^{2\lambda t} \quad \text{as} \quad t \rightarrow \infty \quad (80)$$

So in the course of the expansion of the very late universe, only g_{00} asymptotically vanishes while the space components g_{ii} behave qualitatively in the same manner as the space components of the metric in the Einstein frame $\tilde{g}_{ii}$. Respectively, the asymptotic behavior of the volume measures is as follows:

$$\Phi \approx \frac{b_g V_1 - 2V_2}{V_1} \sqrt{-g} \sim e^{3\lambda t} \quad as \quad t \rightarrow \infty \quad (81)$$

The GES is asymptotically restored that can be seen from the asymptotic time behavior of the conservation law (67)

$$\tilde{\nabla}_\mu J^\mu \sim const \cdot e^{-2\alpha\phi/M_p} \sim \frac{1}{t}. \quad (82)$$

C. The case $b_g V_1 = 2V_2$

In this case the asymptotic form of $V(\phi)$ is

$$V(\phi) \approx \frac{M^8}{2b_g V_1} e^{-4\alpha\phi/M_p} \quad (83)$$

and ζ asymptotically approaches zero according to

$$\zeta = \frac{b_g M^4 e^{-2\alpha\phi/M_p}}{V_1 + M^4 e^{-2\alpha\phi/M_p}} \rightarrow 0 \quad as \quad \phi \rightarrow \infty. \quad (84)$$

Similar to the previous subsection, the analytic form of the asymptotic (as $\phi \gg M_p$) cosmological solution exists for a particular value of the parameter $\alpha = \sqrt{3/32}$:

$$\phi(t) = const. + \frac{M_p}{4\alpha} \ln(M_p t), \quad a(t) \propto t^{1/3} e^{\lambda t}, \quad \lambda = \frac{1}{M_p} \sqrt{\frac{\Lambda}{3}}, \quad (85)$$

where now

$$\Lambda = \frac{V_1}{2b_g} \quad (86)$$

For this solution we obtain the following asymptotic cosmic time behavior for the components of the metric $g_{\mu\nu}$ and volume measures:

$$g_{00} \sim \frac{1}{t^{1/4}} \rightarrow 0; \quad g_{ii} \sim -t^{5/12} e^{2\lambda t} \quad as \quad t \rightarrow \infty \quad (87)$$

$$\sqrt{-g} \sim \sqrt{t} e^{3\lambda t}, \quad \Phi \sim e^{3\lambda t} \quad as \quad t \rightarrow \infty \quad (88)$$

The asymptotic time behavior of the conservation law describing the asymptotic restoration of the GES is the same as in Eq.(82).

D. The case $0 < b_g V_1 < 2V_2$

In this case the potential $U(\phi)$, Eq.(65), has an absolute minimum

$$\Lambda = U(\phi_{min}) = \frac{V_2}{b_g^2} \quad at \quad \phi = \phi_{min} = -\frac{M_p}{2\alpha} \ln \left(\frac{2V_2 - b_g V_1}{b_g M^4} \right). \quad (89)$$

The spatially flat universe described in the Einstein frame with the metric $\tilde{g}_{\mu\nu} = diag(1, -a^2, -a^2, -a^2)$, in a finite time[23] reaches this ground state where it expands exponentially

$$a \propto e^{\lambda t}, \quad \lambda = M_p^{-1} \sqrt{\Lambda/3} \quad (90)$$

and Λ is given by Eq.(89). A surprising feature of this case is that ζ , Eq.(68), disappears in the minimum:

$$\zeta(\phi_{min}) = 0 \quad (91)$$

The components of the metric $g_{\mu\nu}$ in the ground state are as follows

$$g_{00}|_{(ground\ state)} = \left(\frac{2V_2 - b_g V_1}{b_g^3 M^4} \right)^{1/2} = const, \quad g_{ii}|_{(ground\ state)} = -g_{00}|_{(ground\ state)} \cdot e^{2\lambda t} \quad (92)$$

with the respective behavior of the metrical volume measure $\sqrt{-g} \propto \exp(3\lambda t)$. Hence the manifold volume measure in the ground state disappears

$$\Phi|_{(ground\ state)} = 0 \quad (93)$$

in view of Eq.(91).

Disappearance of the manifold volume measure Φ in the ground state may not allow to get the equation (6) by varying the φ_a fields in the action (45). Therefore in the conservation law (67) one should use the current in the form $j^\mu = L_1 B_a^\mu \varphi_a$ as we have noticed after Eq.(40). Recall that L_1 is constituted by the terms of the Lagrangian in (45) coupled to the measure Φ . However, after using the gravitational equation obtained by varying $g^{\mu\nu}$ in (45) and substituting the ground state value $\phi = \phi_{min}$ into L_1 , we obtain $L_1 = M^4$. Hence, the conservation law (67) in the ground state reads just

$$\tilde{\nabla}_\mu J^\mu|_{(ground\ state)} = 0. \quad (94)$$




FIG. 3: The phase portrait (in the phase plane $(\phi, \dot{\phi})$) for the model with $\alpha = 0.2$, $\delta = 0.1$, $V_1 = 10M^4$ and $V_2 = 9.9b_g M^4$. The region with $\rho > 0$ is divided into two dynamically disconnected regions by the line $Q_1(\phi, \dot{\phi}) = 0$. To the left of this line $Q_1 > 0$ (the appropriate zone we call zone 1) and to the right $Q_1 < 0$. The $\rho > 0$ region to the right of the line $Q_1(\phi, \dot{\phi}) = 0$ is divided into two zones (zone 2 and zone 3) by the line $Q_2 = 0$ (the latter coincides with the line where $w = -1$). In zone 2 $w > -1$ but $c_s^2 < 0$. In zone 3 $w < -1$ and $c_s^2 > 0$. Phase curves started in zone 2 cross the line $w = -1$. All phase curves in zone 3 exhibit processes with super-accelerating expansion of the universe. Besides all the phase curves in zone 3 demonstrate dynamical attractor behavior to the line which asymptotically, as $\phi \rightarrow \infty$, approaches the straight line $\dot{\phi} = 0$.




FIG. 4: For the model with $\alpha = 0.2$, $\delta = 0.1$, $V_1 = 10M^4$ and $V_2 = 9.9b_g M^4$: typical scalar factor dependence of ϕ (Fig.(a)) and of the energy density ρ , defined by Eq.(56), (Fig(b)) in the regime corresponding to the phase curves started in zone 2. Both graphs correspond to the initial conditions $\phi_{in} = M_p$, $\dot{\phi}_{in} = 5.7M^2/\sqrt{b_g}$; ρ increases approaching asymptotically the value $\frac{M^4}{b_g} e^{5.52}$




FIG. 5: For the same model and with the same initial conditions as in Fig. 4: crossing the phantom divide $w = -1$ and changing sign of the total volume measure $(\Phi + b_\phi \sqrt{-g})$ in the scalar field ϕ kinetic term (in the underlying action (45)) occur simultaneously.

IX. SIGN INDEFINITENESS OF THE MANIFOLD VOLUME MEASURE AS THE ORIGIN OF A PHANTOM DARK ENERGY

We turn now to the non fine-tuned case of the model of Sec.VII applied to the spatially flat universe. We start from a short review of our recent results[33] concerning qualitative structure of the appropriate dynamical system which

consists of Eq.(58) and the equation

$$\left(\frac{\dot{a}}{a}\right)^2 = \frac{1}{3M_p^2}\rho \quad (95)$$

where the energy density ρ is defined by Eq.(56). The case of the interest of this section is realized when the parameters of the model satisfy the condition

$$(b_g + b_\phi)V_1 - 2V_2 < 0 \quad (96)$$

In this case the phase plane has a very interesting structure presented in Fig.3. Recall that the functions Q_1 , Q_2 , Q_3 are defined by Eqs.(60)-(62).

We are interested in the equation of state $w = p/\rho < -1$, where pressure p and energy density ρ are given by Eqs.(56) and (57). The line indicated in Fig. 3 as "line $w = -1$ " coincides with the line $Q_2(\phi, X) = 0$ because

$$w + 1 = \frac{X}{\rho} \cdot \frac{Q_2}{[b_g(M^4 e^{-2\alpha\phi/M_p} + V_1) - V_2]} \quad (97)$$

Phase curves in zone 3 correspond to the cosmological solutions with the equation of state $w < -1$. In zone 2, $w > -1$ but this zone has no physical meaning since the squared sound speed of perturbations

$$c_s^2 = \frac{Q_2}{Q_1} \quad (98)$$

is negative in zone 3. But in zone 2, $c_s^2 > 0$. Some details of numerical solutions describing the cross of the phantom divide $w = -1$ and the super-accelerating expansion of the universe are presented in Figs. 4 and 5.

Note that the superaccelerating cosmological expansion is obtained here without introducing an explicit phantom scalar field into the underlying action (45). In Ref.[33] we have discussed this effect from the point of view of the effective k-essence model realized in the Einstein frame when starting from the action (45). A deeper analysis of the same effect yields the conclusion that the true and profound *origin of the appearance of an effective phantom dynamics in our model is sign-indefiniteness of the manifold volume measure* Φ . In fact, using the constraint (52), Eqs.(97) and (47) it is easy to show that

$$\Phi + b_\phi\sqrt{-g} = (w + 1) \frac{\rho}{4X} \frac{[M^4 e^{-2\alpha\phi/M_p} + V_1 + \delta \cdot b_g X]}{[b_g(M^4 e^{-2\alpha\phi/M_p} + V_1) - V_2]} a^3, \quad (99)$$

where a is the scale factor. The expression in the l.h.s of this equation is the total volume measure of the ϕ kinetic term in the underlying action (45):

$$\int d^4x (\Phi + b_\phi\sqrt{-g}) \frac{1}{2} g^{\mu\nu} \phi_{,\mu} \phi_{,\nu} \quad (100)$$

The sign of this volume measure coincides with the sign of $w+1$ as well as with the sign of the function Q_2 (see Eq.(97)). In Fig. 5 we present the result of numerical solution for the scale factor dependence of w and $(\Phi + b_\phi\sqrt{-g})/a^3$. Thus *crossing the phantom divide occurs when the total volume measure of the ϕ kinetic term in the underlying action changes sign from positive to negative for dynamical reasons*. This dynamical effect appears here as a dynamically well-founded alternative to the usually postulated phantom kinetic term of a scalar field Lagrangian[41].

X. SUMMARY AND DISCUSSION

Introducing the space-time manifold volume element (2) and adding the appropriate degrees of freedom to a set of traditional variables (metric, connection, matter fields) we reveal that such a two measures theory (TMT) takes up a special position between alternative theories. First, the equations of motion can be rewritten in the Einstein frame (where the space-time becomes Riemannian) with the *same Newtonian constant* as in the underlying action (where the space-time is generically non-Riemannian). Second, the theory possesses remarkable features in what it concerns the CC problem. Third, the TMT model with spontaneously broken dilatation symmetry satisfies all existing tests of GR. There are other interesting results, for example a possibility of a dynamical protection from the initial singularity of the curvature.

In this paper we have studied the behavior of the manifold volume measure Φ and the metric tensor $g_{\mu\nu}$ (used in the underlying TMT action) in cosmological solutions for a number of scalar field models of dark energy. We have made a special accent on the *sign indefiniteness of the manifold volume measure* Φ that may yield interesting physical effects. An example of such type of effects we have seen in Sec.IX: the total volume measure of the dilaton scalar field kinetic term in the underlying action can change sign from positive to negative in the course of dynamical evolution of the late time universe. In the Einstein frame, this transition corresponds to *the crossing of the phantom divide of the dark energy*.

We have found out that in all studied models, the transition to the ground state is *always accompanied by a certain degeneracy* either in the metric (e.g., in g_{00} or in all components) or in the manifold volume measure Φ , or even in both of them. This result differs sharply from what was expected e.g. in Refs.[2]-[4] where degenerate metric solutions have been associated with high curvature and temperature phases. One should only take into account that degeneracy of $g_{\mu\nu}$ and/or Φ in the (transition to) ground state takes place only when one works with the set of variables of the underlying TMT action. In the Einstein frame, we deal with the effective picture where the measure Φ does not present at all and the metric tensor $\tilde{g}_{\mu\nu}$ (see Eqs.(20) or (47)) has the same regularity properties as in GR. The regularity of $\tilde{g}_{\mu\nu}$ results from the singularity of the transformations (20) or (47): degeneracy of $g_{\mu\nu}$ in a discrete set of moments is compensated by a singularity of ζ .

A. The CC problem

TMT provides two different possibilities for resolution of the CC problem: one which guarantees zero CC without fine tuning (see however the end of Sec.VI and Appendix B); another which allows an unexpected way to reach a tiny CC. Which of these possibilities is realized depends on the sign of the integration constant sM^4 , $s = \pm 1$. We are going now to discuss these two issues.

1. The case $\Lambda = 0$ in TMT

This case is of a special interest for two reasons. First, as it was shown earlier[33], the conditions of the Weinberg's no-go theorem[36] fail and a transition to a zero CC state in TMT can be realized without fine tuning. This becomes possible for example if $V_1(\phi) > 0$ and the integration constant $sM^4 < 0$. Second, as we have shown in Sec.V, in the course of transition to a zero CC state, $g_{\mu\nu}$ and Φ oscillate synchronously around zero and they cross zero each time t_i ($i = 1, 2, 3, \dots$) when the scalar field ϕ crosses the (zero) absolute minimum of the potential (27) (or of the potential (65) for the model of Sec.VII with $V_1 < 0$, see [33]).

One should recall that $\zeta(x)$ does not have its own dynamics: its values at the space-time point x are determined directly and immediately by the local configuration of the matter fields and gravity through the algebraic constraint, which is nothing but a consistency condition of equations of motion. $\zeta(x)$ does not possess inertia and therefore it changes together and synchronously with changing matter and gravity fields. This notion is very important when trying to answer the natural question: can oscillations of $\zeta(x)$ be a source for particle creation? The answer is - no, it cannot. In fact, there is a coupling of ζ with fermions. But the structure of this coupling in the Einstein frame has very surprising features which we will shortly review in the next subsection. Here we are only formulating the conclusion: emergence of even a tiny amount of fermionic matter immediately yields a rearrangement of the vacuum[65] in such a way that ζ instantly ceases the regime of oscillations and rapidly enters into a regime of monotonous approach to a nonzero constant. It is interesting to note that the latter effect may explain why the present day cosmological constant most likely is tiny but nonzero, in spite of the existence of a fine tuning free classical solution described a transition to the $\Lambda = 0$ state.

An overall change of sign of $g_{\mu\nu}$ in the course of these oscillations means a change of the signature from $(+ - - -)$ to $(- + + +)$ and vice versa, while oscillations of the sign of Φ describe the change of orientation of the space-time manifold. The latter means that the arena of the gravitational dynamics should contain two space-time manifolds with opposite orientations. The discrete set of changes of the orientations happens in the form of a *smooth* dynamical process in the course of which the space-time passes *the "degenerate" phase where both the metrical structure and the total 4D-volume measure disappear*. The latter means also that the term "orientation of the space-time manifold" loses any sense at moments t_i ($i = 1, 2, 3, \dots$). We conclude therefore that two 4D differentiable manifolds with opposite orientations (described by means of a sign indefinite volume 4-form) equipped with connection and metrical structure still are not enough to describe the arena of the gravitational dynamics: the complete description of the space-time dynamics requires also the mentioned degenerate phase. This situation is somewhat similar to that discussed in Introduction: first-order formulation of GR where the degenerate phase with $g_{\mu\nu} = 0$ should be also added[2],[3],[4]; see e.g. the recent discussion by Bañados[17] where the limiting process $g_{\mu\nu} \rightarrow 0$ is analyzed.

A new interesting feature of ground states in TMT we have revealed in the present paper concerns the so-called global Einstein symmetry (GES), Eqs.(35)-(38), which turns out generically to be explicitly broken in all models with non-trivial dynamics. The surprising result we have discovered here on the basis of a number of models is that *the GES is restored in the course of transitions to the ground state* in all models considered. Hence its subgroup of the sign inversions of $g_{\mu\nu}$ and Φ , Eq.(39), is also restored. Therefore the oscillations of $g_{\mu\nu}$ and Φ around zero in the course of transition to a $\Lambda = 0$ ground state provoke a wish to compare this *dynamical effect* with the attempts to solve the old CC problem developed in Refs.[42]-[44]. The main idea of these approaches is that the field theory or at least the ground state[43] should be invariant under transformations of a discrete symmetry. According to Refs.[42]-[44] it might be either an invariance under the metric reversal symmetry or under the space-time coordinate transformations with the imaginary unit i : $x^A \rightarrow ix^A$. In contrast with these approaches, in TMT there is no need to postulate such exotic enough symmetries. Nevertheless we have seen that sign inversions of $g_{\mu\nu}$ emerge as a dynamical effect in the course of the cosmological evolution and this effect has indeed a relation to the resolution of the old CC problem.

2. The case of a tiny CC

In the scalar field models of dark energy, an interesting feature of TMT consists in a possibility to provide a small value of the CC. If in the model of Sec.VIII.B, the parameter $V_2 < 0$ and $|V_2| \gg b_g V_1$ then the CC can be very small without the need for V_1 and V_2 to be very small. For example, if V_1 is determined by the energy scale of electroweak symmetry breaking $V_1 \sim (10^3 \text{ GeV})^4$ and V_2 is determined by the Planck scale $V_2 \sim (10^{18} \text{ GeV})^4$ then $\Lambda_1 \sim (10^{-3} \text{ eV})^4$. Along with such a seesaw mechanism[25], [45], there exists another way to explain the smallness of the CC applicable in all types of scenarios discussed in Secs.VIII.B-VIII.D (see also Appendix A). As one can see from Eqs.(72), (85) and (89), the value of Λ appears to be inverse proportional[66] to the dimensionless parameter b_g which characterizes the relative strength of the 'manifold' and 'metrical' parts of the gravitational action. If for example $V_1 \sim (10^3 \text{ GeV})^4$ then for getting $\Lambda_1 \sim (10^{-3} \text{ eV})^4$ one should assume that $b_g \sim 10^{60}$. Such a large value of b_g (see Eq.(9)) permits to formulate a *correspondence principle*[33] *between TMT and regular (i.e. one-measure) field theories*: when $\zeta/b_g \ll 1$ then one can neglect the gravitational term in L_1 with respect to that in L_2 (see Eq.(9) or Eq.(17) or Eq.(45)). More detailed analysis shows that in such a case the manifold volume measure $\Phi = \zeta\sqrt{-g}$ has no a dynamical effect and TMT is reduced to GR. This happens e.g. in the model of Sec.VIII.C where the late time evolution proceeds in a quintessence-like manner: the energy density decreases to the cosmological constant, Eq.(86), and $\zeta \rightarrow 0$, Eq.(84). Another example is the model of Sec.VIII.D where $\Phi = 0$ in the ground state, Eq.(91), while $\sqrt{-g}$ is finite. However generically ζ/b_g is not small, as it happens for example in the quintessence-like scenario of the late time universe in the model of Sec.VIII.B (see Eq.(74)).

B. Possibilities for predictions of new physical effects

1. Short review of the TMT model with spontaneously broken dilatation symmetry in the presence of matter

It would be interesting to find out other possible physical manifestations of the sign indefiniteness of the manifold volume measure. In fact, the model with spontaneously broken dilatation symmetry studied in Secs.VII-IX and in Ref.[33] allows extensions which include fermion and gauge fields[29]-[31] or, alternatively, dust as a phenomenological matter model[34]. In the former case, for example, the constraint (52) is modified to the following

$$\frac{1}{(\zeta + b)^2} \left\{ (b - \zeta) \left[M^4 e^{-2\alpha\phi/M_p} + V_1 \right] - 2V_2 \right\} = \frac{\mu(\zeta - \zeta_1)(\zeta - \zeta_2)}{2(\zeta + k)^2(\zeta + b)^{1/2}} \bar{\Psi} \Psi, \quad (101)$$

where Ψ is the fermion field in the Einstein frame and for simplicity we have chosen $\delta = 0$ (that is $b_\phi = b_g = b$); $\zeta_{1,2}$ are defined by

$$\zeta_{1,2} = \frac{1}{2} \left[k - 3h \pm \sqrt{(k - 3h)^2 + 8b(k - h) - 4kh} \right]. \quad (102)$$

and the dimensionless parameters k and h appear in the underlying action in the total volume measures of the fermion kinetic term

$$\int d^4x e^{\alpha\phi/M_p} (\Phi + k\sqrt{-g}) \frac{i}{2} \bar{\Psi} \left(\gamma^a e_a^\mu \vec{\nabla}_\mu - \overleftarrow{\nabla}_\mu \gamma^a e_a^\mu \right) \Psi \quad (103)$$

and the fermion mass term

$$- \int d^4x e^{\frac{3}{2}\alpha\phi/M_p} (\Phi + h\sqrt{-g}) \mu \bar{\Psi} \Psi \quad (104)$$

respectively. Note that the fermion equation in the Einstein frame has a canonical form but the mass of the fermion turns out ζ dependent

$$m(\zeta) = \frac{\mu(\zeta + h)}{(\zeta + k)(\zeta + b)^{1/2}} \quad (105)$$

The constraint (101) describes the local balance between the fermion energy density and the scalar field ϕ contribution to the dark energy density in the space-time region where the wave function of the primordial fermion is not equal to zero. By means of this balance the constraint determines the scalar $\zeta(x)$.

In the case of dust as a phenomenological matter model, the r.h.s. of the constraint (101) looks

$$\frac{\zeta - b_m + 2b}{2\sqrt{\zeta + b}} m \tilde{n}, \quad (106)$$

where the dimensionless parameter b_m appears in the total volume measure of the dust contribution to the underlying action

$$S_m = \int (\Phi + b_m \sqrt{-g}) L_m d^4x \quad (107)$$

$$L_m = -m \sum_i \int e^{\frac{1}{2}\alpha\phi/M_p} \sqrt{g_{\alpha\beta}} \frac{dx_i^\alpha}{d\lambda} \frac{dx_i^\beta}{d\lambda} \frac{\delta^4(x - x_i(\lambda))}{\sqrt{-g}} d\lambda \quad (108)$$

and m is the mass parameter.

The wonderful feature of these models in the Einstein frame consists of the exact coincidence of the following three quantities: a) the noncanonical (in comparison with GR) terms in the energy-momentum tensor; b) the effective coupling "constant" of the dilaton ϕ to the matter (up to the factor α/M_p); c) the expressions in the r.h.s. of the above mentioned constraints (101) and (106) for fermionic matter and dust respectively. For *matter in normal conditions*, the local matter energy density (i.e. in the space-time region occupied by the matter) is many orders of magnitude larger than the vacuum energy density. Detailed analysis[29]-[31],[34] shows that when the matter is in the normal conditions, the balance dictated by the constraint becomes possible if ζ with very high accuracy takes the constant values: $\zeta \approx \zeta_1$ or $\zeta \approx \zeta_2$ for fermions (and therefore the fermion masses become constant)) and $\zeta \approx b_m - 2b$ for the dust. Then the mentioned three quantities simultaneously become extremely small. Besides for the matter in normal conditions the gravitational equations are reduced to the canonical GR equations. The practical disappearance of the dilaton-to-matter coupling "constant" for the matter in normal conditions which occurs without fine tuning of the parameters allows us to assert that in such type of models the fifth force problem is resolved[31],[34].

It does not mean however that matter does not interact with the dilaton at all. When the matter is in states different from normal, the effect of dilaton-to-matter coupling may yield new very interesting phenomena. One of such effects appears when the neutrino energy density decreases to the order of magnitude close to the vacuum energy density. The latter can happen due to spreading of the neutrino wave packet. Then the cold gas of uniformly distributed nonrelativistic neutrinos causes a reconstruction of the vacuum to a state with $\zeta \rightarrow |k|$ and as a result the neutrino gas rapidly transmute into an exotic state called neutrino dark energy(see e.g. Ref.[47]). This effect was studied in details in Ref.[31] where we have shown that transmutation from the pure scalar field dark energy to the neutrino dark energy regime is favorable from the energetic point of view.

2. Prediction of strong gravity effect in high energy physics experiments

For the solutions $\zeta \approx \zeta_1$ or $\zeta \approx \zeta_2$ of the constraint (101), the l.h.s. of the constraint has the order of magnitude close to the vacuum energy density. There exists however another solution if one allows a possibility that in the core of the support of the fermion wave function the local dark energy density may be much bigger than the vacuum energy density. Such a solution turns out to be possible as fermion density is very big and ζ becomes negative and close enough to the value $\zeta \approx -b$. Then the solution of the constraint (101) looks[29]

$$\frac{1}{\sqrt{\zeta + b}} \approx \left[\frac{\mu(b - h)}{4M^4 b(b - k)} \bar{\Psi} \Psi e^{2\alpha\phi/M_p} \right]^{1/3}. \quad (109)$$

In such a case, instead of constant masses, as it was for $\zeta \approx \zeta_{1,2}$, Eq.(105) results in the following fermion self-interaction term in the effective fermion Lagrangian

$$L_{selfint}^{ferm} = 3 \left[\frac{1}{b} \left(\frac{\mu(b-h)}{4M(b-k)} \bar{\Psi}\Psi \right)^4 e^{2\alpha\phi/M_p} \right]^{1/3}. \quad (110)$$

It is very interesting that the described effect is the direct consequence of the strong gravity. In fact, in the regime where $\zeta + b \ll 1$ the effective Newton constant in the gravitational term of underlying action(45)

$$S_{grav} = - \int d^4x \sqrt{-g} \frac{\zeta + b}{\kappa b} R(\Gamma, g) e^{\alpha\phi/M_p} \quad (111)$$

becomes anomalously large. Recall that for simplicity we have chosen here $b_\phi = b_g = b$. But if one do not to imply this fine tuning then one can immediately see from Eqs.(49)-(51) that in the Einstein frame the regime of the strong gravity dictated by the dense fermion matter is manifested for the dilaton too.

The coupling constant in Eq.(110) is dimensionless and depends exponentially of the dilaton ϕ if one can regard ϕ as a background field $\phi = \bar{\phi}$. But in a more general case Eq.(110) may be treated as describing an anomalous dilaton-to-fermion interaction very much different from the discussed above case of interaction of the dilaton to the fermion matter in normal conditions where the coupling constant practically vanishes. Such an anomalous dilaton-to-fermion interaction should result in creation of quanta of the dilaton field in processes with very heavy fermions. The probability of these processes is of course proportional to the Newton constant M_p^{-2} . But the new effect consists of the fact that the effective coupling constant of the anomalous dilaton-to-fermion interaction is proportional to $e^{2\alpha\bar{\phi}/3M_p}$. If the dilaton is the scalar field responsible for the quintessential inflation type of the cosmological scenario[48] then one should expect an exponential amplification of the effective coupling of this interaction in the present day universe in comparison with the early universe. One can hope that the described effect of the strong gravity might be revealed in the LHC experiments in the form of missing energy due to the multiple production of quanta of the dilaton field (recall that coupling of the dilaton to fermions in normal conditions practically vanishes and therefore the dilaton will not be observed after being emitted).

3. Some other possible effects

1. *Dark matter as effect of gravitational enhancement.* In the case of dust as a phenomenological matter model, the constraint (101) with the r.h.s. (106) is the fifth degree algebraic equation with respect to $\sqrt{\zeta + b}$. There are some indications that in a certain region of the parameters a solution of the constraint exists which could provide a very interesting effect of an amplification of the gravitational field of visible diluted galactic and intergalactic dust or/and neutrinos. Such an effect might imply that the dark matter is not a new sort of matter but it is just a result of a so far unknown enhancement of the gravitational field of low density states of usual matter.

2. *Dilaton to photon coupling.* Astrophysical observations of few last years indicate anomalously large transparency of the Universe to gamma rays[49],[50]. It is hard to explain this astrophysical puzzle in the framework of extragalactic background light. Recently a natural mechanism was suggested by De Angelis, Mansutti and Roncadelli[51] in order to resolve this puzzle. The idea is to suppose that there exists a very light spin-zero boson coupled to the photon:

$$L_{\phi\gamma} = -\frac{1}{4\mu} F_{\mu\nu} \tilde{F}_{\mu\nu} \phi. \quad (112)$$

where μ is a mass parameter. In the context of quintessential scenario such a coupling was studied by Carroll[52]. Then $\gamma \longrightarrow \phi \longrightarrow \gamma$ oscillations emerge which explain[51] the observed transparency of the Universe to gamma rays in a natural way if mass of the spin-zero boson $m < 10^{-10} eV$. The crucial feature of this boson is that no other coupling of this scalar to matter exists. In the standard quintessence models this feature seems to be a real problem. But in TMT, as we already mentioned (see also Refs.[31],[34]) the dilaton playing the role of quintessence field decouples from matter in normal conditions. At the same time its coupling to the photon in the form (112) is not suppressed.

3. *Creation of a universe in the laboratory.* A theoretical attempt by Farhi, Guth and Guven to describe a creation of a universe in the laboratory[53] runs across a need to allow vanishing and changing sign of $\sqrt{-g}$. In Ref.[53], this need is naturally regarded as a pathology. If similar approach to the problem of creation of a universe in the laboratory could be formulated in the framework of TMT then instead of $\sqrt{-g}$ there should appear a linear combination of Φ and $\sqrt{-g}$ which, as we already know, is able to vanish and change sign. In recent paper[54] by Guendelman and Sakai a model of child universe production without initial singularities was studied. To provide the desirable absence

of initial singularity a crucial point is that the energy momentum tensor of the domain wall should be dominated by a sort of phantom energy. A possible way to realize this idea is to apply the dynamical brane tension[55] obtained when using the modified volume measure similar to the signed measure Φ of the present paper. So it could be that applying the notions explored in the present paper one can obtain also a framework for formulating non singular child universe production.

4. *Unparticle physics.* ζ dependence of the fermion mass, Eq.(105), together with the constraint (101) can be treated as a $\bar{\Psi}\Psi$ dependence of the fermion mass. This means that in states different from the normal one, the fermion mass spectrum may be continuous, that allows to think of a possibility to establish relation with the idea of unparticle physics[56].

Note finally that for the matter in normal conditions the model does not impose essential constraints on the parameters of the model (such as b_g, b_ϕ, b_m, k, h). But the appropriate constraints should appear when more progress in the study of the listed and another possible new effects will be achieved.

XI. ACKNOWLEDGEMENTS

We acknowledge V. Goldstein, M. Lin, E. Nissimov and S. Pacheva for useful discussions of some mathematical subjects. We also thank M. Duff for explaining us his approach and M. Bañados for useful conversations. We are also grateful to the referee whose constructive remarks assisted us in the improvement of this paper.

APPENDIX A: THE GROUND STATE WITH NON ZERO CC IN MODEL I

Let us consider the scalar field model I (Eqs.(17) and (28)) with $\delta = 0$ where we now choose a positive integration constant ($s = +1$) and the parameters $V_2^{(0)} < 0, b_g\mu_1^2 > \mu_2^2$. Then the ground state is realized for $\phi = 0$ and the vacuum energy is

$$\Lambda = V_{eff}(0) = \frac{M^8}{4b_gM^4 - V_2^{(0)}} \quad (A1)$$

In this ground state, both the measure $\Phi > 0$ and all components of the metric tensor $g_{\mu\nu}$ are regular. Note that the presence of the free dimensionless parameter b_g in the denominator allows again to reach a small vacuum energy by means of the correspondence principle discussed in item 2 of Sec.X.

APPENDIX B: GLOBAL EINSTEIN SYMMETRY DOES NOT GUARANTEE RESOLUTION OF THE CC PROBLEM

In the model (44), the gravitational equations are modified to the following

$$G_{\mu\nu}(\tilde{g}) = \frac{\kappa}{2} \left[\phi_{,\mu}\phi_{,\nu} - \frac{1}{2}\tilde{g}_{\mu\nu}X + \frac{b_g[sM^4 + V_1(\phi) + 2\lambda\zeta] - V_2(\phi) - \lambda\zeta^2}{(\zeta + b_g)^2} \right] \quad (B1)$$

while the form of the scalar field ϕ equation remains the same as in Eq.(25). However the constraint is now very much differs from Eq.(26):

$$4\lambda\zeta^2 + [sM^4 + V_1(\phi) - 2b_g\lambda]\zeta + 2V_2(\phi) - b_g[sM^4 + V_1(\phi)] = 0 \quad (B2)$$

One can see from Eq.(B1) that ζ -dependence emerges now in the numerator of the effective potential. Besides, it is evident that in contrast with what was in Sec.V, the regime with $\zeta \rightarrow \infty$ cannot be a solution of the constraint. It is evident that a zero minimum of the effective potential cannot be now reached without fine tuning. Thus although the second term in the action (44) is invariant under the GES, adding this term we loss the ability to resolve the old CC problem.

- [2] Hawking S W *Nucl.Phys.* 1978 B144 349; Hawking S W 1979 in *Recent Developments in Gravitation* ed M Levy and S Deser (New York; Plenum)
- [3] D'Auria R and Regge T 1982 *Nucl. Phys. B* **195** 308
- [4] Tseytlin A A 1982 *J. Phys. A: Math. Gen.* **15** L105
- [5] Horowitz Gary T 1991 *Class Quantum Grav.* **8** 587
- [6] Ashtekar A 1991 *Lectures on Non-Perturbative Canonical Gravity* (World Scientific)
- [7] Jacobson T and Smolin L 1988 *Nucl. Phys. B* **299** 295
- [8] Dray T, Manogue C A and Tucker R W 1991 *Gen. Rel. Grav.* **23** 967
- [9] Ellis G, Sumeruk A, Coule D and Hellaby C 1992 *Class. Quantum Grav.* **9** 1535
- [10] Elizalde E, Odintsov S and Romeo A 1994 *Class. Quant. Grav.* **11** 61
- [11] Dray T, Ellis G, Hellaby C and Corinne A. Manogue C A 1997 *Gen. Rel. Grav.* **29** 591
- [12] Dray T, Ellis G, Hellaby C 2001 *Gen. Rel. Grav.* **33** 1041
- [13] Borowiec A, Francaviglia M and Volovich I 2007. *Int. J. Geom. Meth. Mod. Phys.* **4** 647
- [14] Mars M, Senovilla Jose M M and Vera R 2008 *Phys. Rev. D* **77** 027501
- [15] Witten E 1988 *Commun. Math. Phys.* **117** 353
Witten E 1988 *Nucl. Phys. B* **311** 46
- [16] Giddings S B 1991 *Physics Letters B* **268** 17
- [17] Bañados M 2007 *Class. Quantum Grav.* **24** 5911
Bañados M 2008 *Phys.Rev.* **D 77** 123534
- [18] Cohn D L, *Measure Theory*, Birkhauser, Boston, 1993.
- [19] Taylor J G 1979 *Phys. Rev.* **19** 2336
- [20] Wilczek F 1998 *Phys. Rev. Lett.* **80** 4851
- [21] Mosna R A and Saa A 2005 *J.Math.Phys.* **46** 112502
- [22] Guendelman E I and Kaganovich A B *Phys. Rev.* 1996 **D53** 7020; *Mod. Phys. Lett.* 1997 **A12** 2421; *Phys. Rev.* **D55** 5970; *Mod. Phys. Lett.* 1997 **A12** 2421; *Phys. Rev.* 1997 **D56** 3548; *Mod. Phys. Lett.* 1998 **A13** 1583.
- [23] Guendelman E I and Kaganovich A B *Phys. Rev.* 1998 **D57** 7200).
- [24] Guendelman E I and Kaganovich A B *Phys. Rev.* 1999 **D60** 065004.
- [25] Guendelman E I 1999 *Mod. Phys. Lett.* **A14**, 1043; *Class. Quant. Grav.* 2000 **17** 361; gr-qc/9906025; *Mod. Phys. Lett.* 1999 **AA4**, 1397; gr-qc/9901067; hep-th/0106085; *Found. Phys.* 2001 **31** 1019;
- [26] Kaganovich A B 2001 *Phys. Rev.* **D63**, 025022.
- [27] Guendelman E I and Katz O 2003 *Class. Quant. Grav.* **20** 1715
- [28] Guendelman E I 1997 *Phys. Lett.* **B412** 42; Guendelman E I 2003 gr-qc/0303048; Guendelman E I and Spallucci E 2003 hep-th/0311102.
- [29] Guendelman E I and Kaganovich A B 2002 *Int. J. Mod. Phys.* **A17** 417.
- [30] Guendelman E I and Kaganovich A B 2002 *Mod. Phys. Lett.* **AA7** 1227 (2002).
- [31] Guendelman E I and Kaganovich A B 2004 hep-th/0411188; *Int.J.Mod.Phys.* 2006 **A21** 4373.
- [32] Guendelman E I and Kaganovich A B 2006 in *Paris 2005, Albert Einstein's century*, AIP Conf.Proc. 2006 861 875, Paris; hep-th/0603229.
- [33] Guendelman E I and Kaganovich A B 2007 *Phys.Rev.* **D75** 083505.
- [34] Guendelman E I and Kaganovich A B 2008 *Annals Phys.* **323** 866.
- [35] Comelli D arXiv:0704.1802 [gr-qc].
- [36] Weinberg S 1989 *Rev. Mod. Phys.* **61** 1
- [37] Unruh W G 1989 *Phys. Rev.* 1989 **D40** 1048.
- [38] Ng Y Jack and van Dam H 1991 *J. Math. Phys.* **32** 1337.
- [39] Chiba T, Okabe T and Yamaguchi M 2000 *Phys.Rev.* **D62** 023511; Armendariz-Picon C, Mukhanov V and Steinhardt P J 2000 *Phys. Rev. Lett.* **85** 4438; *Phys. Rev.* 2001 **D63** 103510; Chiba T 2002 *Phys.Rev.* **D66** 063514.
- [40] Armendariz-Picon C., Damour T and Mukhanov V F 1999 *Phys.Lett.* **B458** 209.
- [41] Caldwell R R *Phys.Lett.* 2002 **B545** 23; Gibbons G W 2003, hep-th/0302199.
- [42] Erdem R 2005 *Phys. Lett.* **B621** 11; *Phys. Lett.* 2006 **B639** 348; *J. Phys.* 2007 **A40** 6945.
- [43] Nobbenhuis S 2006 *Found. Phys.* **36** 613; 't Hooft G, Nobbenhuis S 2006 *Class. Quant. Grav.* **23** 3819.
- [44] Duff M J and Kalkkinen J. 2006 *Nucl. Phys.* **B758** 161; 2007 *Nucl. Phys.* **B760** 64.
- [45] Arkani-Hamed N, Hall L J, Kolda C F and Murayama H 2000 *Phys.Rev.Lett.* **85** 4434.
- [46] R. MacKenzie, F. Wilczek and A. Zee 1984 *Phys. Rev. Lett.* **53** 2203.
- [47] R Fardon, A. E. Nelson, N. Weiner 2004 *JCAP* **0410** 005.
- [48] P.J.E. Peebles and A. Vilenkin 1999 *Phys.Rev* **D59** 063505.
- [49] Uchiyama Y, Aharonian F, Tanaka T, Takahashi T and Maeda 2007 *Nature* **449** 576.
- [50] Mazin D, Raue M 2007 *Astron.Astrophys.* **471** 439.
- [51] De Angelis A, Mansutti O and Roncadelli M 2007 *Phys.Rev.* **D76** 121301.
- [52] Carroll S 1998 *Phys. Rev. Lett.* **81** 3067.
- [53] Farhi E, Guth A H and Guven J 1990 *Nucl. Phys.* **B339** 417.
- [54] Guendelman E I and Sakai N 2008 *Phys. Rev.* **D77** 125002.
- [55] Guendelman E I, Kaganovich A B, Nissimov E, Pacheva S 2002 *Phys. Rev.* **D66** 046003; "Impact of dynamical tensions in modified string and brane theories", Presented at 5th International Workshop on Lie Theory and Its Applications in Physics, Varna, Bulgaria, 16-22 Jun 2003, H.D. Doebner and V. Dobrev Eds., World Scientific, 2004. Published in *Varna*

2003, *Lie theory and its applications in physics V* 241-250 e-Print: hep-th/0401083.

- [56] Georgi H 2007 *Phys. Rev. Lett.* **98** 221601.
- [57] An opposite view on the role of the volume element has been studied by Wilczek[20]. Another idea of modified volume element was studied in Ref.[21].
- [58] For a more detailed discussion of the role of scalars $\varphi_a(x)$ in the TMT dynamics, see the end of Sec.II.
- [59] See also Ref.[33] where we study in detail a model with dilatation symmetry which also results in the k -essence type dynamics
- [60] For the case $s = +1$ and the ground state with nonzero cosmological constant see Appendix A
- [61] Since the metric in the Einstein frame $\tilde{g}_{\mu\nu}$ is diagonal, Eq.(29), it is clear from the transformation (20) that $g_{\mu\nu}$ is also diagonal.
- [62] Note however that these oscillations do not effect the sign of the metrical volume measure $g = \det(g_{\mu\nu})$ used in the underlying action (17). This notion is useful when comparing our model with an approach developed in Refs.[42]-[44]
- [63] for the model of Sec.IV it means that in Eq.(17) we take the limit $b_g \rightarrow 0$, $b_\phi \rightarrow 0$ and $V_2 \rightarrow 0$)
- [64] Note that the pure gravity model of Sec.III is invariant both under the LES and the GES.
- [65] A possibility of a vacuum deformation in a different approach has been shown by MacKenzie, Wilczek and Zee[46]
- [66] In the pure gravity model, Sec.III, Λ is proportional to b_g